Table 3 (South) to Part 660, Subpart F—Non-Trawl Rockfish Conservation Areas

	and Trip Limits for Open Access Gears South of 40°10′ N. lat.									
Та	Table 3 (South) to Part 660, Subpart F – Non-Trawl Rockfish Conservation Areas and Trip Limits for Open Access Gears South of 40°10' N. lat.									
Other limits and requirements apply Read									/16/2021	
		JAN-FEB	MAR-APR	MAY-JUN	JUL-AUG	SEP	-OCT	NOV-DE	С	
Ro	ckfish Conservation Area (RCA) ^{1/} :									
	40 [*] 10' N. lat 38°57.5' N. lat.				e ^{1/} - 125 fm line ^{1/}					
2	38°57.5' N. lat34°27' N. lat.	50 fm line ¹⁷ - 125 fm line ¹⁷								
3	South of 34°27' N. lat.	100 fm line ^{1/} - 150 fm line ^{1/} (also applies around islands)								
See §§660.60 and 660.230 for additional gear, trip limit and conservation area requirements and restrictions. See §§660.70-660.74 and §§660.76-660.79 for								for		
State trip limits and seasons may be more restrictive than Federal trip limits or seasons, particularly in waters off Oregon and California.										
4	Minor Slope Rockfish ^{2/} & Darkblotched rockfish	10,000 lbs/ 2 months, of which no more than 2,500 lbs may be blackgill rockfish								
5	Splitnose rockfish	200 lbs/ month								
	Sablefish									
7	40 10' N. lat 36 00' N. lat.	600 lbs. daily, or 1 landing per week up to 2,000 lbs., not to exceed 4,000 lbs/2 months								
8	South of 36°00' N. lat.	2,000 lbs/week, not to exceed 6,000 lbs/2 months								
9	Shortpine thornyheads									
10	40 [*] 10' N. lat 34 [*] 27' N. lat.	50lb/ month								
	Longspine thornyheads									
12	40 10' N. lat 34 27' N. lat.			50	lb/ month					
	Shortpine thornyheads and longspine									
14	South of 34°27' N. lat.	100 lbs/day, no more than 1,000 lbs/ 2 months								
	Dover sole, arrowtooth flounder,									
	petrale sole, English sole, starry	5,000 lbs/ month								
	flounder, Other Flatfish ^{3/8/}									
	Whiting	300 lbs/ month								
	Minor Shelf Rockfish ^{2/}	4 000 lbs / 0 startle of thickers may then 400 lbs seem by seeming								
20	77 77 78 78 78 78 78 78 78 78 78 78 78 7	4,000 lbs. / 2 months, of which no more than 400 lbs. may be vermillon								
21	South of 34 27' N. lat.	3,000 lbs. / 2 months, of which no more than 1,200lbs. may be vermilion								
	Widow									
23	40 10' N. lat 34 27' N. lat.		6,000 lbs. / 2 months 4,000 lbs. / 2 months							
24	South of 34°27' N. lat.			4,000 1	bs. / 2 months					
	Chilipepper									
26	40 10' N. lat 34 27' N. lat.	6,000 lbs. / 2 months								
27	South of 34 27' N. lat.	4,000 lbs. / 2 months								
28	Shortbelly Rockfish									
29	South of 40°10' N. lat.) lb/ month					
	Canary rockfish				lbs/ 2 months					
	Yelloweye rockfish Cowcod				LOSED					
	Bronzespotted rockfish				LOSED					
	Bocaccio				lbs/ 2 months					
	toactio 4,000 lbs/ 2 months									
31	Shallow nearshore 4/			2 000	lbs/ 2 months					
32	Deeper nearshore 5/				lbs/ 2 months					
	California Scorpionfish	3.500 lbs/ 2 months								
	Lingcod ^{6/}				bs / months				$\overline{}$	
	Pacific cod				lbs/ 2 months					
99				1,000	DO E IIIOIIIII					

150,000 lbs/ 2

Unlimited

Unlimited

Unlimited

100,000 lbs/ 2 months

200,000 lbs/ 2 months

36 Spiny dogfish

38 Big skate

37 Longnose skate

39 Other Fish^{7/} & Cabezon in California